

AD616410



ON THE SEPARATION OF EXPONENTIALS

Richard Bellman  
Mathematics Division  
The RAND Corporation


P-1846

December 11, 1959

Approved for Release by NSA on 09-11-2013 pursuant to E.O. 13526

COPY	1	OF	10
HARD COPY	\$.	1.00	
MICROFICHE	\$.	0.50	

4p



Reproduced by

The RAND Corporation • Santa Monica • California

The views expressed in this paper are not necessarily those of the Corporation

PROCESSING COPY

EVALUATION COPY

ARCHIVE COPY

# SUMMARY

In a number of different fields, it is known as a result of a priori theoretical analysis that a sequence,  $\{u_n\}$ , obtained experimentally, has the representation

$$(1) \quad u_n = \sum_{k=1}^N c_k e^{\lambda_k n}.$$

Even if the dimension,  $N$ , is known, the determination of the parameters  $c_k$  and  $\lambda_k$  can be a matter of some difficulty. If  $N$  is not known initially, the problem is naturally more complex.

The purpose of this note is to outline a method designed to yield the value of  $N$  without the determination of the  $c_k$  and  $\lambda_k$ . In some cases, this is all that is desired.

## ON THE SEPARATION OF EXPONENTIALS

Richard Bellman

1. Introduction

In a number of different fields, it is known as a result of a priori theoretical analysis that a sequence,  $\{u_n\}$ , obtained experimentally, has the representation

$$(1) \quad u_n = \sum_{k=1}^N c_k e^{\lambda_k n}.$$

Even if the dimension,  $N$ , is known, the determination of the parameters  $c_k$  and  $\lambda_k$  can be a matter of some difficulty; cf. the discussions in [1] and [2]. If  $N$  is not known initially, the problem is naturally more complex.

The purpose of this note is to outline a method designed to yield the value of  $N$  without the determination of the  $c_k$  and  $\lambda_k$ . In some cases, this is all that is desired.

2. Recurrence Relations

We start from the frequently used fact that a sequence such as that given above can be generated by means of a recurrence relation of the form

$$(1) \quad v(t + N) = a_1 v(t + N - 1) + \dots + a_N v(t), \quad t = 0, 1, \dots,$$

with  $v(0), v(1), \dots, v(N - 1)$  prescribed. Let  $v(t)$  be a solution of (1), and consider the determinant

$$(2) \quad C_k(t) = \begin{vmatrix} v(t) & v(t + 1) & \dots & v(t + k) \\ v(t + 1) & v(t + 2) & \dots & v(t + k + 1) \\ \vdots & \vdots & \ddots & \vdots \\ v(t + k) & v(t + k + 1) & \dots & v(t + 2k) \end{vmatrix}.$$

Assuming, as we may, that in (1.1)  $c_1 \neq 0$ ,  $i = 1, 2, \dots, N$ , and that  $\lambda_i \neq \lambda_j$ ,  $i \neq j$ , it is easily demonstrated that

- (3) (a)  $c_k(t) = 0$ ,  $k \geq N$ ,  $t = 0, 1, \dots$ ,  
 (b)  $c_{N-1}(t) = c_{N-1}(0)a_N^t \neq 0$ ,  $t = 0, 1, \dots$ ,  
 (c)  $c_k(t) \neq 0$ ,  $k < N - 1$ , for large  $t$ .

These facts can be used to determine  $N$ . Whatever information is available concerning the reality of the  $\lambda_k$  can be used to shorten the work.

### 3. Prediction Theory

Once  $N$  has been determined, we can focus upon the parameters  $a_1$ , and from these obtain the  $c_k$  and  $\lambda_k$ . Perhaps the easiest procedure is to determine the values of the  $a_1$  which minimize

$$(1) \quad Q(a) = \sum_{t=0}^T (u_{t+N} - a_1 u_{t+N-1} - \dots - a_N u_t)^2,$$

for some choice of  $T$ . Since this problem is a discrete version of the basic problem of Wiener prediction theory, (the Kolmogorov version) methods developed in that field (for example, those of Levinson given in the appendix to Wiener's book) can be used.

### 4. Discussion

The problem is actually more complex, since what is measured experimentally is not  $u_n$ , but  $w_n = u_n + r_n$ , where  $r_n$  is a random quantity. It may well be that in this case, a reasonable procedure is to minimize the expected value of the quadratic form appearing in (3.1), and to choose a value of  $N$  which minimizes this.

1. D. C. Gardner, J. C. Gardner, G. Laush, and W. W. Meinke, "Method for the analysis of multicomponent exponential decay curves," J. Chem. Phys., vol. 31, 1959, pp. 978-986.
2. C. Lanczos, Applied Analysis, Prentice-Hall, 1956.